

WE CLAIM:

-1-

A vaccine for providing passive immunity to *Sarcocystis neurona* infection comprising antibodies which are against at least one epitope of a unique 16 ( $\pm 4$ ) or 30 ( $\pm 4$ ) antigen of *Sarcocystis neurona*.

-2-

The vaccine of Claim 1 wherein the antibodies are selected from the group consisting of polyclonal antibodies and monoclonal antibodies.

-3-

The vaccine of claim 1 wherein the vaccine is provided in a pharmaceutically accepted carrier.

-4-

A vaccine for active immunization of an equid against a *Sarcocystis neurona* infection comprising at least one epitope of a unique 16 ( $\pm 4$ ) or 30 ( $\pm 4$ ) antigen of *Sarcocystis neurona*.

-5-

The vaccine of Claim 4 wherein the antigen is a recombinant polypeptide produced in a plasmid in a microorganism other than *Sarcocystis neurona*.

-6-

The vaccine of Claim 5 wherein the microorganism is an *E. coli*.

-46-

-7-

5

The vaccine of Claim 6 wherein the antigen is a fusion polypeptide wherein an amino end or a carboxyl end of the antigen is fused to all or a portion of a polypeptide that facilitates isolation of the antigen from the microorganism in which the antigen is produced.

-8-

The vaccine of Claim 7 wherein the polypeptide is selected from the group consisting of glutathione S-transferase, protein A, maltose binding protein, and polyhistidine.

-9-

The vaccine of Claim 6 wherein the vaccine is provided in a pharmaceutically accepted carrier.

-10-

A vaccine for protecting an equid from a *Sarcocystis neurona* infection comprising a DNA that encodes at least one epitope of a 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen of *Sarcocystis neurona*.

-11-

The vaccine of Claim 10 wherein the DNA is operably linked to a promoter to enable transcription of the DNA in a cell of an equid.

-12-

The vaccine of Claim 10 wherein the vaccine is provided in a pharmaceutically accepted carrier.

0382130686-02200000

A method for vaccinating an equid against a *Sarcocystis neurona* infection comprising:

- 5 (a) providing a recombinant antigen of *Sarcocystis neurona* produced from a microorganism culture wherein the microorganism contains a DNA that encodes at least one epitope of a 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen of *Sarcocystis neurona*; and  
(b) vaccinating the equid.

The method of Claim 13 wherein the recombinant antigen is in a pharmaceutically accepted carrier.

5 The method of Claim 13 wherein the recombinant antigen is a fusion polypeptide which is fused at the amino terminus or carboxyl terminus to a polypeptide that facilitates the isolation of the recombinant antigen.

The method of Claim 15 wherein the polypeptide includes all or a portion of the polypeptide selected from the group consisting of glutathione S-transferase, protein A, maltose binding protein, and polyhistidine.

The method of Claim 15 wherein the DNA is in a plasmid in a microorganism wherein the DNA is operably linked to a promoter which enables transcription of the DNA to produce the recombinant antigen for the vaccine.

A method for vaccinating an equid against a *Sarcocystis neurona* infection comprising:

(a) providing in a carrier solution a DNA in a plasmid which encodes at least one epitope of a 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen of *Sarcocystis neurona*; and

5 (b) vaccinating the equid with the DNA in the carrier solution.

The method of Claim 18 wherein the carrier solution is a saline solution.

The method of Claim 18 wherein the DNA is operably linked to a promoter to enable transcription of the DNA in a cell of the equid.

A method for providing passive immunity to a *Sarcocystis neurona* infection in an equid comprising:

(a) providing antibodies against at least one epitope of a 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen of *Sarcocystis neurona* wherein the antibodies are selected from the group consisting of polyclonal antibodies and monoclonal antibodies; and

5 (b) inoculating the equid.

The method of Claim 21 wherein the antibodies are provided in a pharmaceutically accepted carrier.

*SAC*  
*AZ*

5

10

0000000000 - 622406

-49-

-23-

A method for producing a polypeptide comprising:

- (a) providing a microorganism in a culture containing a DNA encoding a fusion polypeptide comprising at least one epitope of a 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen of *Sarcocystis neurona* and a polypeptide that facilitates isolation of the fusion polypeptide;
- (b) culturing the microorganism in a culture to produce the fusion polypeptide; and
- (c) isolating the fusion polypeptide.

-24-

The method of Claim 23 wherein isolating the fusion polypeptide is by affinity chromatography.

-25-

The method of Claim 24 wherein the polypeptide is all or a portion of protein A and the affinity chromatography comprises an IgG-linked resin.

-26-

The method of Claim 24 wherein the polypeptide is polyhistidine and the affinity chromatography comprises a  $Ni^{2+}$  resin.

-27-

The method of Claim 24 wherein the polypeptide is glutathione S-transferase and the affinity chromatography comprises a glutathione Sepharose 4B resin.

-50-

-28-

The method of ~~Claim 24~~ wherein the polypeptide is maltose binding protein and the affinity chromatography comprises an amylose resin.

-29-

A method for producing an antibody comprising:

- (a) providing a microorganism in a culture containing a DNA encoding a fusion polypeptide comprising at least one epitope of a 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen of *Sarcocystis neurona* and a polypeptide that facilitates isolation of the fusion polypeptide;
- (b) culturing the microorganism in a culture to produce the fusion polypeptide;
- (c) isolating the fusion polypeptide;
- (d) producing the antibody from the polypeptide.

30-

A method for producing a monoclonal antibody comprising:

- (a) providing a microorganism in a culture containing a DNA encoding a fusion polypeptide comprising at least one epitope of a 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen of *Sarcocystis neurona* and a polypeptide that facilitates isolation of the fusion polypeptide;
- (b) culturing the microorganism in a culture to produce the fusion polypeptide;
- (c) isolating the fusion polypeptide;
- (d) producing the monoclonal antibody from the polypeptide.

-51-

-31-

The method of Claim 29 or 30 wherein isolating the fusion polypeptide is by affinity chromatography.

-32-

The method of Claim 31 wherein the polypeptide is all or a portion of protein A and the affinity chromatography comprises an IgG-linked resin.

-33-

The method of Claim 31 wherein the polypeptide is polyhistidine and the affinity chromatography comprises a  $\text{Ni}^{2+}$  resin.

-34-

The method of Claim 31 wherein the polypeptide is glutathione S-transferase and the affinity chromatography comprises a glutathione Sepharose 4B resin.

-35-

The method of Claim 31 wherein the polypeptide is maltose binding protein and the affinity chromatography comprises an amylose resin.

-36-

A monoclonal antibody that selectively binds to a 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen of *Sarcocystis neurona*.

-37-

An isolated recombinant protein encoded by a cDNA produced from RNA of *Sarcocystis neurona* encoding a 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen.

-52-

-38-

An isolated DNA that encodes a 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen of *Sarcocystis neurona*.

-39-

A bacterial clone containing a plasmid comprising a DNA encoding a 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen of *Sarcocystis neurona*.

-40-

The bacterial clone of Claim 39 wherein the clone expresses the 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen of *Sarcocystis neurona*.

-41-

A vaccine for an equid comprising an isolated recombinant protein encoded by a cDNA produced from mRNA of *Sarcocystis neurona* encoding a protein which is a 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen, and a vaccine carrier.

5

-42-

A vaccine for an equid comprising a recombinant virus vector containing DNA encoding a 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen of *Sarcocystis neurona*, and a vaccine carrier.

-43-

The vaccine of Claim 42 wherein the recombinant virus is selected from the group consisting of equid herpesvirus, vaccinia virus, canary poxvirus, raccoon poxvirus, and adenovirus.

-53-

-44-

A DNA vaccine for an equid comprising a plasmid containing DNA encoding a 16 ( $\pm 4$ ) and/or 30 ( $\pm 4$ ) kDa protein of *Sarcocystis neurona*.

-45-

A method for protecting an equid against *Sarcocystis neurona* which comprises providing a vaccine that when injected into the equid causes the equid to produce antibodies against a 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen of the *Sarcocystis neurona* wherein the antibodies prevent infection by the *Sarcocystis neurona*.

5

-46-

*July 23* The method of Claim 45 wherein the vaccine comprises the 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen in a vaccine carrier.

-47-

The method of Claim 45 wherein the vaccine is a recombinant virus vector that expresses the 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen.

-48-

The method of Claim 47 wherein the recombinant virus vector is selected from the group consisting of equine herpesvirus, vaccinia virus, canary poxvirus, raccoon poxvirus, and adenovirus.

-49-

The method of Claim 45 wherein the vaccine comprises a DNA plasmid encoding the 16 ( $\pm 4$ ) kDa antigen and/or 30 ( $\pm 4$ ) kDa antigen.

055012666 - 022666

-54-

-50-

*Sub esp*  
The method of ~~Claim 45~~ wherein the vaccine is  
administered by a vaccination route selected from the  
group consisting of intranasal administration,  
intramuscular injection, intraperitoneal injection,  
intradermal injection, and subcutaneous injection.

5